ASEAN Biological Threats Surveillance Centre

Situation Report of Ebola Disease Outbreak caused by the Bundibugyo virus (BDBV) in the Democratic Republic of the Congo

November 13, 2026

With Support by:

Situation at a Glance

- On May 17, 2026, the World Health Organization (WHO) officially declared the Ebola disease outbreak caused by the Bundibugyo virus (BDBV) in the Democratic Republic of the Congo (DRC) a Public Health Emergency of International Concern (PHEIC) under the International Health Regulations (2005).
- **On August 14**, WHO reassessed the outbreak risk and maintained its previous assessment: very high in the DRC, high in countries sharing land borders with the DRC, and low in the rest of the African Region and globally.
- **As of November 11**, a total of 11,197 cases with 5,471 deaths **(CFR: 48.9%)** have been confirmed in DRC across 67 health zones in eight provinces. A total of 3,045 patients recovered, representing a recovery rate of 27.2%.




Figure 1. Map of countries reporting confirmed cases of BDBV as of November 11, 2026

- Since the start of the outbreak, **six countries have reported imported BVD cases linked to affected areas**, highlighting the risk of international spread. **Uganda, France, Germany, and the UK** reported cases associated with travel to the DRC, while **Kenya** reported imported cases linked to travel from the DRC via Uganda. **Nigeria** subsequently reported an imported case in a traveler returning from Uganda.

Situation Update

Democratic Republic of the Congo

- **On May 5**, 2026, WHO received an alert regarding a high-mortality outbreak in Mongbwalu Health Zone, Ituri Province, including four healthcare worker deaths within four days. Retrospective investigations identified the earliest suspected case as a healthcare worker who developed fever, bleeding, vomiting, and severe malaise on 24 April and later died in Bunia.

- **On May 15,** Laboratory testing confirmed Bundibugyo virus disease (BVD), and the Ministry of Health officially declared the country's 17th Ebola outbreak. At the time, 246 suspected cases and 80 deaths had been reported across Rwampara, Mongbwalu, and Bunia health zones.
- **On May 24**, the outbreak expanded to 11 health zones across Ituri, North Kivu, and South Kivu, with 105 confirmed cases and 10 deaths (CFR: 9.5%), alongside 906 suspected cases and 223 suspected deaths.
- **Between June 14 to 20,** the country reported 174 new cases and 66 deaths (CFR: 37.9%), bringing cumulative totals to 1,094 confirmed cases and 277 deaths.
- **Between July 9 and October 16**, the outbreak expanded beyond the initial affected provinces, with **Tshopo** and **Haut-Uélé** added on 9 July, followed by **Bas-Uélé** (12 August), **Sud-Ubangi** (10 September), and **Nord-Ubangi** (16 October), increasing the number of affected provinces from three to eight.
- **By November 11,** DRC confirmed a total of 11,197 cases and 5,471 deaths (CFR=48.9%) from 68 health zones in eight provinces.

Table 1. Distribution of Confirmed BDBV cases in the Democratic Republic of the Congo (DRC), as of November 11, 2026

Province	Health Zones Affected	Cases	Deaths	CFR (%)
Ituri	28	7,765	3,667	47.2
Nord-Kivu	17	2,802	1,547	55.2
Haut-Uele	8	504	218	43.3
Tshopo	7	98	32	32.7
Sud-Kivu	1	3	1	33.3
Bas Uele	4	13	4	30.8
Sud-Ubangi	1	2	1	50.0
Nord-Ubangi	2	10	1	10.0
Total	**68**	**11,197**	**5,471**	**48.9**

Source: Institut National De Sante Publique DRC

The BVD outbreak in the DRC continued to be characterized by sustained transmission and high mortality. Weekly reported cases increased from 8 cases in epidemiological week (EW) 19 to a peak of 856 cases in EW29, reflecting rapid expansion of the outbreak across affected provinces (Figure 2). Although case counts subsequently declined, transmission remained substantial, with weekly totals largely exceeding 375 cases from EW30 onward. Cases decreased from 509 in EW42 to 503 in EW43 and 453 in EW44, suggesting a modest downward trend. In the current reporting period, 277 cases had been reported in EW45, covering November 8–11, indicating continuous transmission.

Mortality trends followed a similar pattern. Weekly deaths increased from 4 deaths in EW19 to a peak of 475 deaths in EW29, before declining in subsequent weeks. Despite this decline, mortality remained persistently high, with weekly deaths ranging from 179 to 365 between EW30 and EW44. As of November 10, DRC had reported 110 deaths in the incomplete EW45 reporting period. The weekly case fatality ratio (CFR) fluctuated throughout the outbreak, remaining above 45% since EW30 and peaking at 56.1% in EW32, indicating continued high mortality among reported cases.

Overall, although weekly case counts and deaths have declined from the peaks observed in late July and early August, transmission remains active across affected provinces. The continued reporting of hundreds of new cases and deaths each week indicates that the outbreak remains a significant public health threat. The apparent decline observed in EW45 should be interpreted cautiously, as the reporting period covers only four days (Figure 2).




Figure 2. Weekly new cases of BVD in the DRC (May 10 – November 11, 2026)
(Source: Institut National De Sante Publique DRC)

Ituri Province remains the most affected province, accounting for 69.3% of all confirmed cases reported to date. The highest proportions of confirmed cases were reported from Bunia (18.5%), Rwampara (10.9%), Nizi (7.8%), and Nizi (7.8%). Together, these three health zones accounted for approximately 37.2% of all confirmed cases, highlighting the concentration of transmission in a limited number of persistent hotspots.

Outside Ituri, North Kivu accounted for 25.0% of confirmed cases, followed by Haut-Uélé (4.5%). The remaining affected provinces contributed relatively small proportions of cases, each accounting for 0.1%–0.9% of the cumulative total, indicating that transmission remains concentrated in a limited number of provinces and health zones.




Figure 3. Distribution of Health Zones affected, as of November 11, 2026

Meanwhile, cases reported from other provinces indicate that transmission is not confined to Ituri alone and that the risk of further spread persists. Continued population movement, challenges in accessing healthcare services, and delays in detection may contribute to ongoing transmission and the emergence of new affected areas.

Uganda

- **On May 11, 2026,** Uganda reported its first confirmed case of BVD in a traveler from the DRC. Laboratory confirmation was obtained on 15 May, and the patient subsequently died. The last confirmed case was reported on 21 June 2026.
- **By July 28, 2026,** Uganda officially **declared the end of the Ebola outbreak** with no further transmission detected. Overall, the outbreak resulted in 20 confirmed cases and two deaths (CFR: 10.0%), with 18 patients recovering (90.0%).

France

- **On June 24, 2026,** France reported an imported laboratory-confirmed BDBV case in a physician returning from a humanitarian mission in Ituri Province, DRC. The patient was isolated after self-reporting symptoms upon arrival on June 23. PCR testing confirmed infection, and five contacts identified from the flight were placed under monitoring. No secondary cases have been reported.

Germany

- **On July 13, 2026,** an imported BVD case involving a United States humanitarian worker was medically evacuated from the DRC to Germany after testing positive for BDBV. At the time of reporting, no evidence of local transmission or secondary cases had been reported in Germany.

Kenya

- **On October 6, 2026**, Kenya reported its first imported BDBV case in a traveler returning from the DRC via Uganda, who died on October 5. A second epidemiologically linked case was confirmed in a household contact under quarantine on October 14.
- **As of October 23**, Kenya reported nine confirmed cases, including two deaths, with all cases linked to the initial importation event. There was no evidence of wider community transmission.
- **By November 11,** Kenya's last Ebola patient has been discharged, with all 135 contacts completing monitoring and no community transmission recorded. The 42-day countdown towards declaring Kenya Ebola-free begins on November 12.

Nigeria

- **On October 20, 2026**, Nigeria reported its first imported BDBV case in a Swiss national with recent travel history to Uganda. On October 26, Switzerland reported the medical evacuation of the patient, who remains in isolation at Geneva University Hospital. Her spouse has since been discharged from hospital. No information on the patient's clinical condition has been released due to data protection requirements, and no additional cases have been reported in Nigeria.

The United Kingdom

- **On November 5, 2026,** the United Kingdom reported the evacuation of a healthcare worker who tested positive for Ebola disease while supporting outbreak response activities in the DRC. The patient was reported to be in stable condition, and UK authorities assessed the risk to the public as low.

Epidemiology

- Bundibugyo virus disease (BVD) is a severe Ebola disease caused by the Bundibugyo virus, an Orthoebolavirus. The current event is the 17th Ebola outbreak reported in the DRC since 1976.
- The outbreak has now surpassed DRC's 2018–2020 outbreak in deaths and is the largest Ebola outbreak ever reported in DRC.

Response Operations

Democratic Republic of the Congo

- Response activities continued across affected provinces. Surveillance at points of entry and points of control (PoE/PoC) remained active, while RCCE interventions supported community awareness, contact tracing, and outbreak response efforts.
- Logistics operations continued to support the response through personnel training, the delivery of nutrition and IPC supplies, and fuel distribution.
- Following confirmation of the second BDBV case on 14 October 2026, Kenya intensified contact tracing and quarantine measures. Of the 128 contacts identified from the index case, 113 had been traced, including 108 contacts placed under quarantine and daily monitoring in Kenya, while relevant authorities were notified for five contacts located abroad. Efforts to locate the remaining contacts continue.
- On October 15, 2026, WHO reaffirmed its support to Kenya's Ebola response following confirmation of a second BDBV case. Response efforts have focused on surveillance, contact tracing, case management, and infection prevention and control (IPC).

United Kingdom

- The British Government arranged the patient's medical evacuation from the DRC to the UK for specialist treatment at a high-consequence infectious disease (HCID) unit, with transfer by specialist ambulance upon arrival. British authorities stated that the National Health Service (NHS) has procedures in place to manage suspected and confirmed Ebola infections.

The WHO Recommendations

- On August 24, 2026, following the second meeting of the IHR Emergency Committee, WHO reaffirmed that the BVD outbreak continues to constitute a PHEIC and issued updated Temporary Recommendations for affected and at-risk countries.
- WHO stressed the importance of community trust, engagement with community leaders and local networks, and acceptance of public health measures, including early detection, case referral, contact tracing, and safe burial practices.
- WHO highlighted the importance of community trust and cross-border collaboration through strengthened surveillance, information sharing, and preparedness for imported cases, while continuing to advise against travel or trade restrictions, flight suspensions, or border closures.
- WHO also underscored the need to accelerate research on BVD vaccines and therapeutics while maintaining core outbreak-control measures, including surveillance, contact tracing, clinical care, IPC, and community engagement.

Risk to the ASEAN Region

Currently, there are no direct (non-stop) flights from the outbreak source airport (Kinshasa – DRC) to all ASEAN Member States (AMS). This allows for significant reduction on immediate importation risk, as passenger volume is diluted across connecting routes, and additional screening opportunities exist at transit hubs. Despite no direct flights, all ASEAN-bound travel requires 1-stop connections, primarily through major global hubs. Further details on flight connectivity can be accessed in the Situation Report of Ebola Disease Outbreak in The Democratic Republic of the Congo and Uganda - 18 May 2026.

References

1. Bluedot (2026) BlueDot Alerts: Ebola in Democratic Republic of the Congo
2. Bluedot (2026) BlueDot Alerts: Ebola in Kenya
3. Bluedot (2026) BlueDot Alerts: Ebola in Nigeria
4. Bluedot (2026) BlueDot Alerts: Ebola in United Kingdom
5. Institut National De Sante Publique (2026). *SitRep N°181/MVE-BDBV/11/11/2026*. Available from: https://insp.cd/sitrep-n181-mve-bdbv-11-11-2026/
6. Ministry of Health of the Republic of Kenya (2026, November 3) *Ebola Response Update. Quarantine complete and start of 42-day countdown*. Available from https://web.facebook.com/MinstryofHealthKE
7. Swiss Info (November 5, 2026) *The husband of a Swiss Ebola patient has been discharged from Zurich University Hospital (USZ)*. Available from https://www.swissinfo.ch/eng/various/husband-of-swiss-ebola-patient-discharged-from-hospital/92334864
8. UK Government (November 4, 2026) *UK citizen returned from DRC for specialist Ebola care*. Available from https://www.gov.uk/government/news/item/24-08-2026-second-meeting-of-the-ihr-emergency-committee-on-the-epidemic-of-ebola-bundibugyo-virus-disease-in-the-democratic-republic-of-the-congo-temporary-recommendations
9. World Health Organization (2026). *BUNDIBUGYO VIRUS DISEASE OUTBREAK Democratic Republic of the Congo | Kenya | Nigeria - Weekly External Situation Report 25, Data as of 7 November 2026.* Available from: https://www.afro.who.int/countries/kenya/publication/bundibugyo-virus-disease-outbreak-democratic-republic-congo-kenya-nigeria-weekly-external-situation
10. World Health Organization (2026, November 6). *Ebola disease caused by Bundibugyo virus - Democratic Republic of the Congo, Kenya and Nigeria*. Available from: https://www.who.int/emergencies/disease-outbreak-news/item/2026-DON625

ASEAN BIOLOGICAL THREATS SURVEILLANCE CENTRE

ASEAN Biological Threat Surveillance Centre (ABVC)
Health Policy Agency
Ministry of Health of Indonesia
Building 6 - Global Health Strategy and Governance Policy, 2nd floor
Jalan Percetakan Negara No. 29 Johar Baru, Jakarta Pusat, Indonesia 10560